WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:		(11) International Publication Number: WO 97/47289
A61K 9/50	A1	(43) International Publication Date: 18 December 1997 (18.12.97)
(21) International Application Number: PCT/US (22) International Filing Date: 9 June 1997 (BE, CH. DE, DK. ES, EI, ER, GR. GR. IE IT IN AGO
(30) Priority Data: 08/662,870 12 June 1996 (12.06.96)	ι	Published With international search report.
(71) Applicant: E-L MANAGEMENT CORPORATION 767 Fifth Avenue, New York, NY 10153 (US).	lus/us	S];
(72) Inventors: KNIGHT, Althea; 308 Herrick Avenue, NJ 07666 (US). ZECCHINO, Julius; 30 Auryanse Closter, NJ 07624 (US). SCHNITTGER, Ste Willetts Place, Huntington Station, NY 11746 (US)	en Cour	erri d
(74) Agents: FLINTOFT, Gerald, J. et al.; Pennie & Edmor 1155 Avenue of the Americas, New York, NY 100	nds LLI 136 (US	P. (5).
·		
(54) Title: STABILIZED HINOKITIOL AND COMPOSIT		CO. T

(57) Abstract

The present invention relates to cosmetic or pharmaceutical compositions containing a preservative-effective amount of hinokitiol-containing microcapsules, the microcapsules comprising a matrix containing collagen and a glycosaminoglycan.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

	Albania	ES	Spain	เร	Lesotho	SI	Slovenia
AL		FI	Finland	LT	Lithuania	SK	Słovakia
AM	Armenia	FR	France	LU	Luxembourg	SN	Senegal
AT	Austria	GA	Gabon	1.V	Letvia	SZ	Swaziland
ΑU	Australia	GB	United Kingdom	MC	Monaco	TD	Chad
AZ	Azerbaijan		_	MD	Republic of Moldova	TG	Togo
BA	Bosnia and Herzegovina	GE	Georgia	MG	Madagascar	TJ	Tajikistan
BB	Barbados	GH	Ghana	MK	The former Yugoslav	TM	Turkmenistan
BE	Belgium	GN	Guinea	MIR	Republic of Macedonia	TR	Turkey
BF	Burkina Faso	GR	Greece		-	TT	Trinidad and Tobago
BG	Bulgaria	HU	Hungary	ML	Mali	ÜA	Ukraine
BJ	Benin	IE	ireland	MN	Mongolia	UG	Uganda
BR	Brazil	IL	Israel	MR	Mauritania	US	United States of America
BY	Belarus	IS	lceland	MW	Malawi	UZ	Uzbekistan
CA	- Canada	IT	lta ly	MX	Mexico		Viet Nam
CF	Central African Republic	JP	Japan	NE	Niger	VN	
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Kores	PL.	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
cz	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	Li	Liechtenstein	SD	Sudan		
	Denmark	LK	Sri Lanka	SE	Sweden		
DK		LR	Liberia	SG	Singapore		
EE	Estonia	LA	2.00.1.2		- 0 :		
i							

WO 97/47289 PCT/US97/09835

STABILIZED HINOKITIOL AND COMPOSITIONS CONTAINING SAME

Field of the Invention

5

The present invention relates to preservatives useful in pharmaceutical and cosmetic compositions. In particular, the invention relates to a stabilized, anti-irritant preservative, hinokitiol.

Background of the Invention

Contamination of pharmaceutical and cosmetic compositions by bacteria and/or fungi can present a serious problem to the formulator. To combat this problem, and in order to prolong shelf life of the product, such compositions frequently contain preservatives which have antimicrobial activity. Given the diverse chemical nature of the available preservatives, however, and the equally diverse nature of the compositions in which they are intended to be used, the ability to successfully use a given preservative in a particular formulation cannot be assumed.

One such preservative is the substance known as 20 hinokitiol. [2-hydroxy-4-(1-methylethyl)-2,4,6cycloheptatrien-1-one; isopropyltropolone]. Hinokitiol is a wood extract obtainable from pine, cedar, etc. Methods for its production are described, for example, in JP 0586396 and JP 05105898. This material is known to have potent antimicrobial 25 effects, and, as such, has been previously used for a variety of purposes. For example, hinokitiol has been used in food preservation and prevention of discoloration (JP 78035145; JP 6277019; JP 6153788; JP 6090661; JP 5344861; JP 59085279); in combination with zinc, or with ethyl alcohol and propylene 30 glycol for antisepsis and/or the treatment of infection(WO 9317559; JP 3133334); in detergent compositions (WO 9205240); and as a wood preservative (JP 1038203). Hinokitiol has also

15

20

25

30

been used in various types of cosmetic and pharmaceutical compositions. For example, DE 4202964 discloses a water-phase hair and body treatment comprising hinokitiol and heliotropin or a cosmetic alcohol, wherein the heliotropin and alcohol is said to synergistically enhance the antimicrobial activity of hinokitiol. Similarly, JP 2243607 describes the use in cosmetic compositions of a preservative containing hinokitiol in combination with phenoxyethanol and phthalate ester. In addition, as shown herein, hinokitiol also possesses a heretofore unappreciated anti-irritancy effect.

Although hinokitiol has been shown to be an effective preservative under a wide variety of conditions, it is not itself completely stable under all conditions in which its use might be desired. For example, its use in compositions in polyethylene plastic containers can result in a discolored It also may not be stable and/or unpleasant-smelling product. after direct contact with light or air. These are limitations which under ordinary circumstances would not be acceptable, and which would recommend against its use in a cosmetic composition, where appearance and fragrance can be critical. Therefore, in order to fully exploit its useful properties as a preservative, there remains a need for a means by which hinokitiol can itself be stabilized in the desired composition, and yet retain its useful biological properties, such as antimicrobial and anti-irritant activity. The present invention provides the means by which hinokitiol can be effectively used in cosmetic compositions.

SUMMARY OF THE INVENTION

The present invention provides cosmetic and pharmaceutical compositions containing hinokitiol-containing microcapsules, the microcapsules comprising a matrix containing collagen and a glycosaminoglycan. The microcapsule provides adequate

stability to the hinokitiol, preventing it from discoloration and unpleasant odor in the composition containing same, while nonetheless permitting the hinokitiol to exert its preservative and anti-irritant effect on the composition containing it. The invention also provides a method for preserving a cosmetic or pharmaceutical composition, comprising adding to the composition a preservative-effective amount of the hinokitiol containing microcapsules, as well as a method for reducing irritation on the skin by application of a cosmetic composition containing an effective amount of hinokitiol-containing microspheres.

DETAILED DESCRIPTION OF THE INVENTION

The microcapsules used in the present invention are prepared according to the methodology disclosed in US Patent No. 5,395,620, the contents of which are incorporated herein by 15 reference. Briefly, the microcapsules are prepared by combining a solution containing atelocollagen and a solution containing a polyholoside, preferably a glycosaminoglycan, in the presence of the hinokitiol and a crosslinking agent. Atelocollagen is a type of collagen from which the telopeptides which crosslink typical collagen have been removed. glycosaminoglycans are well known in the art, and for the present purposes may be selected, for example, from chondroitin 4-sulfate, chondroitin 6-sulfate, dermatan sulfate, heparan sulfate, keratan sulfate, heparin and derivatives thereof; preferably the cosmetically or pharmaceutically acceptable salts, such as calcium or sodium salts, are employed. amount of glycosaminoglycan present in the microcapsules will typically be about 15-50% by weight of the atelocollagen used in the formulation. In absolute terms, the amount of 30 atelocollagen in the starting solution is approximately 0.5-2% by weight, and the glycosaminoglycan solution contains about 0.5-4%, preferably 0.5-2% by weight of glycosaminoglycans.

20

a particularly preferred embodiment, both the atelocollagen and the glycosaminoglycans are of marine origin. Cross-linking reagents useful in preparing the microcapsules include, but are not limited to, an acid dichloride, an acid anhydride, or a dibasic or polybasic carboxylic acid. With respect to the hinokitiol, this material is present in an amount of between 0.01-10% by weight, preferably 0.5-4% by weight, of the starting solution.

The microcapsules containing hinokitiol provide an unexpectedly high level of preservative efficacy, while itself remaining stable in the compositions in which it is included. The retention of antimicrobial activity, at the levels seen herein, is particularly surprising in view of the fact that the hinokitiol itself is not expected to be in direct contact with the medium is intended to preserve. The microcapsules in which the hinokitiol is contained are essentially impermeable, i.e., there is apparently substantially no leakage of the hinokitiol into the medium in which the microcapsules are contained, as evidenced by the absence of unpleasant odor or color in the products containing them. Since the hinokitiol is used in quantities essentially the same as used in its unencapsulated form, the substantially complete retention of antimicrobial activity by the "protected" hinokitiol is quite unexpected.

The encapsulated hinokitiol can be used in a wide variety of cosmetic and/or pharmaceutical compositions, as well as a preservative in food compositions. In particular, the hinokitiol can be used in virtually any form in which such compositions may be formulated, e.g., solutions, colloidal dispersions, emulsions, suspensions, creams, lotions, gels, foams, mousses and the like. Methods and components for preparing such compositions are well known in the art, and can be found for example, in CTFA International Cosmetics Ingredients Dictionary, The Cosmetic, Toiletry, and Fragrance

30

15

20

Association, Washington, DC, 1991. The amount of the hinokitiol-containing microspheres will vary depending on the nature of the composition and the amount of hinokitiol contained in the microspheres, but a preservative-effective amount of hinokitiol per se will generally be in the range of from about 0.1-1% by weight of the composition as a whole, given the amount of hinokitiol in the microspheres is as stated above. It is well within the skill of the art to determine the desired microsphere concentration in the formulation, given these parameters.

The compositions of the present invention will also comprise a pharmaceutically or cosmetically acceptable carrier, in an amount appropriate to accommodate the other components of the formulation. The term "pharmaceutically or cosmetically acceptable carrier" refers to a vehicle, for either pharmaceutical or cosmetic use, which vehicle delivers the active components to the intended target and which will not cause harm to humans or other recipient organisms. As used herein, "pharmaceutical" or "cosmetic" will be understood to encompass both human and animal pharmaceuticals or cosmetics. The carrier may be in any form appropriate to the mode of delivery, for example, solutions, colloidal dispersions, emulsions, suspensions, creams, lotions, gels, foams, mousses and the like.

In a preferred embodiment, the carrier is a suspension, dispersion or emulsion. The emulsion may be an oil-in-water emulsion, or a water-in-oil emulsion. These emulsions contain one or more oil components, an aqueous component, and a specific emulsifier component chosen depending on the nature of the desired emulsion.

The oil component may be any pharmaceutically or cosmetically acceptable material which is substantially insoluble in water. These materials can be found for example in

the CTFA International Dictionary of Cosmetic Ingredients as well as the U.S. Pharmacopoeia or equivalent sources. Suitable oil components include, but are not limited to, natural oils, such as coconut oil; hydrocarbons, such as mineral oil and hydrogenated polyisobutene; fatty alcohols, such as octyldodecanol; esters, such as C12-15 alkyl benzoate; diesters, such as propylene glycol dipelargonate; triesters, such as glyceryl trioctanoate; sterol derivatives, such as lanolin; animal waxes, such as beeswax; plant waxes, such as carnauba; mineral waxes, such as ozokerite; petroleum waxes, 10 such as paraffin wax; synthetic waxes, such as polyethylene; and mixtures thereof. Suitable oil components may also be silicones including, but not limited to, volatile silicones such as cyclomethicone; polymeric silicones such as dimethicone; alkylated derivatives of polymeric silicones, such as cetyl dimethicone and lauryl trimethicone; hydroxylated derivatives of polymeric silicones, such as dimethiconol; and mixtures thereof. The aqueous component refers to any pharmaceutically or cosmetically acceptable material consisting essentially or predominantly of water. 20

For preparation of an oil-in water emulsion, the oil-in-water emulsifier will be an emulsifier having a hydrophilic-lipophilic balance (HLB) of at least 6, or a mixture of such emulsifiers with one or more water-in-oil emulsifiers (i.e., emulsifiers having an HLB of from about 2 to about 6), in which case the type and amount of each emulsifier present in the mixture is selected such that the effective HLB of the resultant oil-in-water emulsifier component is at least about 6. Techniques for combining and ascertaining the effective HLB of a mixture of emulsifiers are known; see L.M.Prince, in M.G. DeNavarre, "The Chemistry and Manufacture of Cosmetics", Volume III, Second Ed., (Continental Press, Orlando, 1975), pp. 25-37.

25

30

10

15

20

30

Suitable oil-in-water emulsifiers include, but are not limited to, sorbitol derivatives, such as sorbitan monolaurate and polysorbate 20; ethoxylated alcohols such as laureth-23, ethoxylated fatty acids such as PEG-1000 stearate; amidoamine derivatives such as stearamidoethyl diethylamine; sulfate esters such as sodium lauryl sulfate; phosphate esters such as DEA cetyl phosphate; fatty acid amine salts such as TEA stearate; and mixtures thereof.

The emulsion may also be a water-in-oil emulsion. For this purpose, a water-in-oil emulsifier is employed. This refers to any cosmetically acceptable emulsifier having an HLB of no greater than 6, preferably from about 2 to about 4.

Suitable water-in-oil emulsifiers include, but are not limited to, sorbitan derivatives such as sorbitan laurate and sorbitan palmitate; alkoxylated alcohols such as laureth-4; hydroxylated derivatives of polymeric silicones, such as dimethicone copolyol; alkylated derivatives of hydroxylated polymeric silicones, such as cetyl dimethicone copolyol; glyceryl esters such as polyglyceryl-4 isostearate; beeswax derivatives such as sodium isostearoyl-2 lactylate; lecithin; and mixtures thereof.

The active component of the pharmaceutical or cosmetic compositions will depend on the intended purpose of the compositions. Examples of such active agents which may form part of the composition include, but are not limited to, those that improve or eradicate age spots, keratoses and wrinkles, analgesics, anesthetics, anti-acne agents, antibacterials, antiyeast agents, antifungal agents, antiviral agents, antidandruff agents, antidermatitis agents, antipruritic agents, antiemetics, antimotion sickness agents, anti-dry skin agents, antiperspirants, antipsoriatic agents, antiseborrheic agents, hair conditioners and hair treatment agents, antiaging

agents, antiwrinkle agents, antiasthmatic agents and bronchodilators, sunscreen agents, antihistamine agents, skin lightening agents, depigmenting agents, vitamins, corticosteroids, tanning agents, hormones, retinoids, topical cardiovascular agents, clotrimazole, ketoconazole, miconozole, griseofulvin, hydroxyzine, diphenhydramine, pramoxine, lidocaine, procaine, mepivacaine, monobenzone, erythromycin, tetracycline, clindamycin, meclocyline, hydroquinone, minocycline, naproxen, ibuprofen, theophylline, cromolyn, albuterol, retinoic acid, 13-cis retinoic acid, hydrocortisone, 10 hydrocortisone 21-acetate, hydrocortisone 17-valerate, hydrocortisone 17-butyrate, betamethasone valerate, betamethasone diproprionate, triamcinolone acetonide, fluocinonide, clobetasol, proprionate, benzoyl peroxide, 15 crotamiton, propranolol, promethazine, vitamin A palmitate, vitamin E acetate and mixtures thereof.

In one preferred embodiment, the composition is a sunscreen-containing composition. The term "sunscreen" as used herein refers to any material which is capable of protecting human skin from ultraviolet radiation having a wavelength of from about 280 to about 400nm, by effectively absorbing such radiation, and/or reflecting or scattering such radiation away from the surface of human skin. Suitable sunscreens for purposes of this invention include but not limited to, titanium derivatives such as titanium dioxide, especially titanium dioxide having an average particle size of from 10 to 100 nanometers, most especially titanium dioxide having an average particle size of from 10 to 100 nanometers and comprising a hydrophobic coating agent to minimize agglomeration; zinc derivatives such as zinc oxide, especially zinc oxide having an average particle size of from 10 to 100 nanometers; melanin and melanin derivatives; dioxybenzone; 2-ethylhexyl 2-cyano-3,3diphenylacrylate; 2-ethylhexyl p-methoxycinnamate; 2-ethylhexyl

20

25

30

salicylate; homosalate; menthyl anthranilate; oxybenzone; octyl dimethyl PABA; red petrolatum; ferulic acid ester; and mixtures thereof.

In addition to those components specifically noted above, the compositions may also comprise additional preservatives, fragrances, emollients, antiseptics, antiinflammatories, antibacterials, stabilizers, antioxidants, vitamins, pigments, dyes, humectants, and propellants, as well as other classes of materials the presence of which in the compositions may be cosmetically, medicinally, or otherwise desired. Such components can be found in the CTFA International Cosmetics Ingredients Dictionary, supra.

The present invention is further illustrated by the following non-limiting examples.

15 EXAMPLE I

Thalaspheres® (30µm) containing hinokitiol are provided by Bioetica, Inc., Portland, Maine). The composition of the Thalaspheres®, in grams per 100g, is as follows:

20	Ingredient	Total	Support phase	Capsules
		,		composition
	Deionized water	81.494	77.86%	89.98%
	Butylene glycol	15.0	21.43%	
	Hinokitiol	2.4		8.0%
25	Xanthan gum	0.5	0.71%	
	atelocollagen	0.441		1.47%
	chondroitin	0.165		0.55%
		100 g	100%	100%
30	equivalent(in g/100	g of produc	t) 70g	30g

SUBSTITUTE SHEET (RULE 26)

BNSDOCID: <WO____9747289A1_1 >

EXAMPLE 2

The encapsulated hinokitiol, as described in Example 1, is incorporated by art-recognized procedures into a sunscreencontaining dispersion, a moisturizing cream emulsion, and a skin desensitizing lotion emulsion. The components are as follows:

SUNSCREEN FORMULATION

	Component	Weight %
10	cyclomethicone/dimethicone copolyol	16.0
	Titanium dioxide Dispersion	13.0
	cyclomethicone	15.5
	dimethicone	10.5
	phenyltrimethicone	6.0
15	hydrogenated lecithin	0.5
	deionized water	36.0
	magnesium sulfate	1.5
	encapsulated hinokitiol	1.0

20 MOIS	STURIZING FORMULATION
---------	-----------------------

	Component	Weight %
	Deionized water	62.0
25	oat stearate	0.9
	Disodium EDTA	0.1
	squalane	5.0
30	cetearyl alcohol/cetearyl glucoside	5.0
	shea butter	6.0
35	sucrose	2.0
	isostearyl neopentanoate	2.0
	cyclomethicone	7.5
40	carbomer	0.25

	glycerine USP 95%	2.5
5	triethanolamine 99%	0.05
•	sodium hyaluronate	2.5
	encapsulated hinokitiol	4.2
10		

	SKIN DESENSITIZING FORMULA	NOITA
	Satin Finish*	Weight %
15	Disodium EDTA	0.1
	Shea butter	0.5
20	rosemary extract	0.1
	glycerine USP 95%	4.5
	gorgonian extract	0.1
25	aluminum starch octenyl succinate	1.2
	carbomer	0.2
30	sucrose	1.8
	caffeine powder	0.4
	triethanolamine 99%	0.05
35	magnesium aluminum sulfate	0.25
	deionized water	36.6

*a commercial (Collaborative Labs) emulsion comprising phenyl trimethicone, cyclomethicone, phosophoglycerides, dimethiconol, phenoxyethanol, carbomer, and triethanolamine

4.2

45

40

encapsulated hinokitiol

The formulations so prepared exhibit no deterioration, no discoloration and no unpleasant odor even after prolonged

storage, showing the encapsulated hinokitiol retains its preservative effect while at the same time being stabilized by the encapsulation.

5 EXAMPLE 3

10

Hinokitiol alone is tested for its ability to prevent irritation. Hinokitiol (unencapsulated, 0.5% by weight) is applied to the ventral forearms of panelists. The material is allowed to absorb for twenty minutes and then Balsam of Peru, an irritant, is applied to the test sites. Skin irritation is measured in terms of increase in skin redness. Cola nitida(10% hydro-alcohol 1:1) is used as a positive control.

The degree of redness is measured with the Minolta chromameter and compared with the positive and negative controls. The positive control is the color of skin treated with Balsam of Peru alone and the negative control is the skin treated with cola material challenged as with the hinokitiol. Hinokitiol exhibits a 71% activity, as compared with an average of 67% for cola.

The encapsulated hinokitiol is then tested for its retained anti-irritant activity. Encapsulated hinokitiol, prepared as described above, is prepared in 4.2% and 2.8% aqueous solution. These solutions are tested against empty microcapsules, as well as against cola nitida(10% hydro-alcohol 1:1), using the same protocol as described in the previous example. Results show that the activity of the encapsulated hinokitiol at 4.2% is approximately equivalent to that of 10% cola (61% vs 64%, respectively) in reducing the onset of irritation due to Balsam of Peru. The lower concentration of hinokitiol capsules shows 53% activity, while empty microcapsules show 28% activity.

What we claim is:

- 1. A cosmetic or pharmaceutical composition containing a preservative-effective amount of hinokitiol-containing microcapsules, the microcapsules comprising a matrix containing collagen and a glycosaminoglycan.
- 2. The composition of claim 1, in which the matrix comprises atelocollagen and chondroitin sulfate.

10

- 3. The composition of claim 1 in which the hinokitiol is present in the microcapsules in an amount of from 0.1-10% by weight.
- 15 4. The composition of claim 1 in which the microcapsules are present in an amount of from about 1-10%.
 - 5. The composition of claim 1 which is in the form of a solution, colloidal dispersion, emulsion, suspension, cream, lotion, gel, foam, or mousse.
 - 6. The composition of claim 5 which is a dispersion.
 - 7. The composition of claim 5 which is an emulsion.

25

30

20

- 8. The composition of claim 7 which is a water-in-oil emulsion.
- 9. The composition of claim 6 which also contains a sunscreen.
 - 10. The composition of claim 9 in which the sunscreen is titanium dioxide.

- 11. The composition of claim 10 in which the microcapsules are present in an amount of about 1-5%.
- 12. A food composition containing a preservative-effective amount of hinokitiol-containing microcapsules, the microcapsules comprising a matrix containing collagen and a glycosaminoglycan.
- 10 13. A method for preserving a pharmaceutical, food or cosmetic composition comprising adding to the composition a preservative effective amount of hinokitiol-containing microcapsules.
- 14. A method for reducing the irritating effects of a cosmetic or pharmaceutical composition on the skin comprising adding to the composition an anti-irritant effective amount of hinokitiol-containing microcapsules.

INTERNATIONAL SEARCH REPORT

International application No. PCT/US97/09835

A 635	LOCATION TO A STATE OF THE STAT			
A. CLA	ASSIFICATION OF SUBJECT MATTER :A61K 9/50			
US CL :424/456				
According	to International Patent Classification (IPC) or to b	oth national classification and IPC		
B. FIE	LDS SEARCHED			
Minimum o	documentation searched (classification system follo	wed by classification symbols)		
U.S. :	424/456			
Documenta	tion searched other than minimum documentation to	the extent that such documents are included	in the fields searched	
Electronic o	data base consulted during the international search	(name of data base and, where practicable	, search terms used)	
C. DOC	UMENTS CONSIDERED TO BE RELEVANT			
Category*				
	Citation of document, with indication, where	appropriate, of the relevant passages	Relevant to claim No.	
Y	US 5,268,174 A (SAKUMA ET A entire document.	L.) 07 December 1993, see	1-14	
(US 5,395,620 A (HUC ET AL.) document.	07 March 1995, see entire	1-14	
l				
- [·			
ŀ				
		1		
Furthe	er documents are listed in the continuation of Box	C. See patent family annex.		
	is categories of cited documents:	"I" later document published after the inter	national filing date or priority	
A* docs to be	ment defining the general state of the art which is not considered to fracticular relevance	date and not in conflict with the applicat principle or theory underlying the inver-	ion but cited to understand the	
	er document published on or after the international filing date	"X" document of particular relevance; the	claimed invention cannot be	
CACO	ment which may throw doubts on priority claim(s) or which is to establish the publication date of another citation or other isl reason (as specified)	considered novel or cannot be considered when the document is taken alone 'Y' document of particular relevance; the	claimed invention cannot be	
O° docu mea	ment referring to an oral disclosure, use, exhibition or other	considered to involve an inventive a combined with one or more other such being obvious to a person skilled in the	decuments such combination	
P* document published prior to the international filing date but later than the priority date claimed document member of the same patent family				
Pate of the actual completion of the international search Date of mailing of the international search report				
18 AUGUST 1997 2 9 AUG 1997				
ame and mailing address of the ISA/US Commissioner of Patents and Trademarks Authorized officer				
Box PCT Washington, D.C. 20231 Raj Bawa, Ph.D.				
csimile No.		Telephone No. (703) 308-2351		
m PCT/ISA	/210 (second sheet)(July 1992)*			